

CD68 [D4B9C] - 156Gd


Catalog: 715601

Clone: D4B9C


Isotype: Rabbit IgG

Reactivity: Human*


Application: MIBI-FFPE

Storage: Supplied in antibody stabilizer with 0.05% sodium azide. Store at 4°C



IHC: CD68 antibody staining of FFPE human tonsil



MIBI: CD68 antibody staining (cyan) of FFPE human tonsil, counterstained with histone H3 (magenta)



IHC: CD68 antibody staining of FFPE human liver



MIBI: CD68 antibody staining (cyan) of FFPE human liver, counterstained with histone H3 (magenta)

Background

CD68 is a marker predominantly expressed on monocytes and macrophages and is used to identify tumor-associated macrophages (TAMs). Macrophages home to specific targets in part through the interaction between CD68 and tissue- and organ-specific lectins or selectins. CD68 staining can be combined with HLA-DR to identify M1 macrophages or with CD163 to identify M2 macrophages, macrophage phenotypes with prognostic associations to outcomes in certain cancers.

Validation

Each lot of conjugated antibody is quality control tested by staining tissue following the MIBI Staining Protocol optimized for the applicable tissue format with subsequent MIBIScope analysis using the appropriate positive and negative tissue field of views. These results are pathologist verified.

Recommended Usage

Human FFPE: 1:100 dilution. For optimal results, the antibody should be titrated for each desired application.

References

Mei, J. et al. Prognostic impact of tumor-associated macrophage infiltration in non-small cell lung cancer: A systemic review and meta-analysis. *Oncotarget*. 2016; **7**(23): 34217-28.

* Conjugate tested on human tissue.